PATIENT NAME:	DATE OF ASSESSMENT:
PATIENT DATE OF BIRTH:	MEDICAL RECORD NUMBER:

Clinical Opioid Withdrawal Score (COWS)

COWS is not required prior to starting buprenorphine, but it can help ensure the patient is ready for the first dose. For each item, write in the number that best describes the patient's signs or symptoms. Rate only the apparent relationship to opiate withdrawal. For example: If heart rate is increased because the patient was jogging just prior to assessment, the increased pulse rate would not add to the score.

Enter scores at time zero, 30 minutes after fir	rst dose, 2 hours after first dose, etc.	Time:	Time:	Time:	Time:
Resting Pulse Rate: Record beats per minute after	r patient is sitting or lying down for one minute				
• 0 - pulse rate 80 or below • 1 - pulse rate 81–100	 2 - pulse rate 101–120 4 - pulse rate greater than 120 				
Sweating: Over past ½ hour not accounted for by	room temperature or activity				
 0 - no chills or flushing 1 - subjective chills or flushing 2 - flushed or observable moistness on face 	3 - beads of sweat on brow or face4 - sweat streaming off face				
Restlessness: Observation during assessment • 0 - able to sit still	• 3 - frequent shifting or extraneous movement of legs/arms				
• 1 - reports difficulty sitting still, but is able to do so	• 5 - unable to sit still for more than a few seconds				
Pupil size					
0 - pupils pinned or normal size for light1 - pupils possibly larger than normal for light	2 - pupils moderately dilated5 - pupils dilated that only rim of the iris is visible				
Bone or joint aches : If patient was having pain preattributed to opiate withdrawal is scored	eviously, only the additional component				
 0 - not present 1 - mild/diffuse discomfort 2 - patient reports severe diffuse aching of joints/muscles 	• 4 - patient is rubbing joints or muscles and is unable to sit still because of discomfort				
Runny nose or tearing: Not accounted for by cold	symptoms or allergy				
0 - none present1 - nasal stuffiness or unusually moist eyes	 2 - nose running or tearing 4 - nose constantly running or tears streaming down cheeks 				
Gl upset: Over last ½ hour	• 2 - nausea or loose stool				
0 - no Gl symptoms1 - stomach cramps	 3 - vomiting or diarrhea 5 - multiple episodes of diarrhea or vomiting				
Tremor: Observation of outstretched hands					
0 - no tremor1 - tremor can be felt, but not observed	2 - slight tremor observable4 - gross tremor or muscle twitching				
Yawning: Observation during assessment olimits 0 - no yawning to 1 - yawning once or twice during assessment	 2 - yawning three or more times during assessment 4 - yawning several times/minute 				
Anxiety or irritability	• 2 - patient obviously irritable or anxious				
 0 - none 1 - patient reports increasing irritability or anxiousness 	4 - patient so irritable or anxious that participation in the assessment is difficult				
Gooseflesh skin	• 3 - piloerection of skin can be felt or hairs standing up on arms				
• 0 - skin is smooth	• 5 - prominent piloerection				
Suggested score to start buprenorphine					
in the ED is COWS \geq 8	TOTAL				
In the ED is COVVS 28	OBSERVER INITIALS				