



CHAPTER 73

Dementia and the Rural Emergency Department

CHRISTOPHER R. CARPENTER,
MD, MSc, FACEP, FAAEM, AGSF¹

Institutional Affiliations:

1. Associate Professor of
Emergency Medicine, Director
of Evidence Based Medicine,
Washington University in St.
Louis, St. Louis, MO

Keywords: rural emergency
medicine, dementia

Vignette

A 78-year-old farmer presents to your rural emergency department (ED) with his family for evaluation of exertional chest pain. While initiating your diagnostic evaluation for chest pain (see Chapter 25) his son who resides out-of-town notes that “Dad seems more forgetful than last year. Last night Mom told me that he cannot find the keys to the truck and he forgot his oldest grandchild’s name this morning.” After carefully inquiring about any recent standing level falls or other trauma, medication changes, or acute illnesses, you contemplate the differential diagnosis of altered mental status (see Chapter 20) and consider the possibility of dementia.

The key questions you contemplate are:

1. *How likely is dementia amongst geriatric patients in the ED?*
2. *What are the clinical implications for my ED patient if I do or do not assess for dementia during an emergency care encounter?*
3. *How do I screen for dementia in the ED?*
4. *What do I do with a patient with abnormal dementia screening test results?*

Epidemiology of Dementia in Rural Settings

Because of aging baby-boomers and the successes of modern medicine prolonging individual’s anticipated lifespan, geriatric adults continue to present to EDs in unprecedented numbers.¹ However, the contemporary ED practice model was not designed for geriatric care.² Consequently, ED healthcare providers rarely screen for geriatric syndromes such as cognitive dysfunction, standing level falls, and baseline functional status.³ Octogenarians with cognitive dysfunction are a potentially vulnerable subset of patients and many reside at home without a reliable safety net.⁴ Multiple studies over the last two decades demonstrate that ED providers fail to recognize cognitive dysfunction in up to 70% of cases.⁵⁻⁸ Outpatient physicians also commonly miss the diagnosis of cognitive dysfunction.^{9,10}

Cognitive dysfunction includes mild cognitive impairment, delirium, and dementia. We review altered mental status (Chapter 20) and delirium (Chapter 80) elsewhere so this chapter will focus on the diagnosis and management of dementia in rural settings. Dementia progresses from mild cognitive impairment to severe, debilitating dementia.¹¹ The most common cause of dementia is Alzheimer’s Disease, but vascular disease, head trauma, and nutritional deficiencies also cause dementia.¹² Ongoing debate questions the value of non-specialty based

dementia screening¹³⁻¹⁵ with opponents noting the lack of valid screening instruments and inadequate curative therapy for this chronic, debilitating neurodegenerative disorder. These biases against dementia detection are a form of therapeutic nihilism and represent real-world barriers to detecting dementia.

Even without a cure for dementia, definitive diagnosis provides emergency clinicians, patients and caregivers several potential benefits. First, emergency care providers establish a baseline mental status in the medical record so that future evaluations can more promptly identify altered mental status as a deviation from baseline. Second, healthcare providers who recognize abnormal mental status earlier can re-evaluate the chief complaint and seek additional history for family and friends to ensure that the true reason for the current ED evaluation has been identified. Third, patients and families with concerns for memory deficits have the otherwise ignored opportunity to seek definitive testing and obtain access to ongoing clinical trials for dementia care. Fourth, early diagnosis provides patients with preclinical dementia with the opportunity to establish and communicate decisions about advanced directives and financial commitments with loved ones. Ideally, early stage dementia is identified by primary care providers, but this is the exception rather than the norm.⁹

When formally assessed, approximately 35% of ED adults over age 65 have cognitive dysfunction as defined by a Mini

Mental Status Exam (MMSE) score of less than 24.^{8,16} Although no rural ED studies have yet assessed the prevalence or impact of dementia, it is common in rural settings worldwide.^{17,18} In fact, patients with dementia may be more common in rural settings than in urban environments.¹⁹ More severe cognitive impairment amongst rural patients is found with increasing age, less formal education, and impaired functional ability.²⁰ However, rural patients and communities often lack awareness of dementia as a geriatric syndrome limiting impetus to seek medical care.²¹ Rural primary care providers are less likely to diagnose dementia compared with their urban counterparts.²² Consequently, rural dementia patients are less likely to receive neurologist or psychiatrist specialty care.²³ Rural nursing residents are more likely to reside in facilities without accreditations or special care program access, which may promote more frequent transfers to the rural ED for evaluation.²⁴

Economic and Clinical Relevance of Dementia to Rural ED Healthcare Providers

Worldwide, over 34 million individuals have dementia, including over 5 million Americans.^{25,26} Dementia care costs \$150 billion annually in the United States.²⁷ These expenses include medical visits, pharmaceutical therapy, diagnostic tests, home nursing resources, and institutionalization, as well as non-medical costs such as consumables and professional and non-professional supervision. These costs are often borne by the patient's family.²⁸ The cost of dementia care worldwide was estimated at \$422 billion, a 34% increase between 2005 and 2009.²⁵ By 2050, the worldwide prevalence of dementia will increase four-fold to about 107 million individuals.²⁹ It is projected that if 30 years from now dementia interventions can delay cognitive decline by just one-year then over 9 million less individuals would require more expensive, higher levels of

care so the impetus is on the healthcare system to continue to identify dementia patients and develop effective management strategies.²⁹ The presence of dementia also affects individual patients and healthcare systems. Dementia is independently associated with short-term ED readmissions, functional decline, institutionalization, and increased hospital length of stay.³⁰⁻³²

Screening for Dementia in the Rural ED

Geriatric patients do not present to the ED with the chief complaint that "I have dementia". Instead, they (or their caregivers on their behalf) present with distinct constellations of symptoms, impairments, and concerns that the astute physician molds into a diagnosis using the art and science of medicine.³³ Dementia will only be identified in the ED by proactive case finding, but the screening does not require a physician's expertise. Physician extenders, nurses, students, or volunteers can screen for dementia using validated, ED-appropriate instruments.³

Traditional instruments to assess for dementia in the clinical setting require time, equipment, training, and personnel that are not readily available in most EDs. Examples of these lengthy instruments include the MMSE and the Montreal Cognitive Assessment instruments.^{34,35} Brief dementia screening instruments for rural environments have been derived, but await validation in ED settings where acute illness-related distress and discomfort, as well as the chaotic milieu, can affect cognitive assessment diagnostic test performance.^{36,37} The ideal ED screening instrument (for dementia or anything else) would be brief (< 2 minutes to administer), sufficiently accurate to rule-in (positive likelihood ratio >10) or rule-out (negative likelihood ratio < 0.1) dementia, easy to remember without requiring special equipment, and acceptable to patients. Unfortunately, this dementia-screening instrument does not exist.³⁸ However, geriatric emergency medicine researchers recently validated several instruments with negative likelihood ratios ≤ 0.10 (Table 1).^{8,16} We will present four of these ED-appropriate screening

TABLE 1: Diagnostic Test Characteristics of ED-Validated Dementia Screening Instruments

	Sensitivity % (95% CI)	Specificity % (95% CI)	Positive Likelihood Ratio (95% CI)	Negative Likelihood Ratio (95% CI)
SBT	95 (88-98)	65 (61-67)	2.7 (2.2-3.0)	0.08 (0.03-0.2)
BAS	95 (88-98)	52 (48-54)	2.0 (1.7-2.2)	0.10 (0.03-0.3)
O3DY	95 (85-99)	51 (46-53)	2.0 (1.6-2.1)	0.10 (0.03-0.3)
cAD8	83 (71-91)	63 (55-68)	2.2 (1.6-2.8)	0.27 (0.1-0.5)

Abbreviations: Confidence Interval (CI), Short Blessed Test (SBT), Brief Alzheimer's Screen (BAS), caregiver-administered AD8 (cAD8).

instruments beginning with the most feasible, least complex and progressing to the more time-consuming, more accurate instruments. We propose the algorithm in **Figure 1** as one pragmatic approach for cognitive dysfunction screening in the rural ED.

The Ottawa 3DY (**Figure 2**) is the easiest instrument to use. It consists of 4 simple questions that the healthcare provider or designated screener can ask while the patient awaits ED evaluation.³⁹ Any incorrect response is an “abnormal” Ottawa 3DY result and merits additional testing, usually outside of

FIGURE 1: Rural ED Dementia Screening Protocol




FIGURE 2: Ottawa 3DY

- | | | |
|-----------------------------------|----------------|-------------|
| 1. <u>What day is today?</u> | Correct | Incorrect |
| 2. <u>What is the date?</u> | Correct | Incorrect |
| 3. <u>Spell “world” backwards</u> | Number Correct | 0 1 2 3 4 5 |
| 4. <u>What year is this?</u> | Correct | Incorrect |

A single incorrect response on any of these four items is consistent with cognitive impairment

the ED. The Short Blessed Test (Figure 3) is the next most feasible dementia-screening instrument, consisting of six basic questions and cognitive tasks.⁴⁰ Patient’s responses are noted and screeners then score the Short Blessed Test based upon

an algebraic weighting formula. The Brief Alzheimer’s Screen (Figure 4) consists of four questions and cognitive tasks with scoring based upon an algebraic interpretation of correct and incorrect responses.⁴¹ The AD8 (Figure 5) consists of eight

FIGURE 3: Short Blessed Test

Instructions to the patient: Now I would like to ask you some questions to check your memory and concentration. Some of them may be easy and some of them may be hard.

	Correct	Incorrect
1. What year is it now?	0	1
2. What month is this?	0	1

Please repeat this name and address after me:

John Brown, 42 Market Street, Chicago; John Brown, 42 Market Street, Chicago; John Brown, 42 Market Street, Chicago
(underline words repeated correctly in each trial)

Trials to learn _____ (if unable to do in 3 trials = C)

3. Without looking at your watch or clock, tell me what time it is. (If response is vague, prompt for specific response within 1-hour) hat year is it now?

Correct	Incorrect
0	1

4. Count aloud backwards from 20 to 1 (mark correctly sequenced numerals—if subject starts counting forward or forgets the task, repeat instructions and score one error)

0 1 2 Errors

20 19 18 17 16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1

5. Say the months of the year in reverse order. If the tester needs to prompt with the last name of the month of the year, one error should be scored—mark correctly sequenced months.

0 1 2 Errors

D N O S A JL JN MY AP MR F J

6. Repeat the name and address you were asked to remember.
 (John Brown, 42 Market Street, Chicago)

0 1 2 Errors

_____, _____, _____, _____, _____, _____

Item	Errors	Weighting Factor	Final Item Score
1		x4	
2		x3	
3		x3	
4		x2	
5		x2	
6		x2	

<p>Sum Total (range 0-28)</p> <p>0-4 = normal cognition</p> <p>5-9 = questionable impairment</p> <p>≥ 10 = impairment consistent with dementia</p>

FIGURE 4: Brief Alzheimer's Screen

Instructions to the patient: I would like to ask you some questions that ask you to use your memory. I am going to name three objects. Please wait until I say all three words, then repeat them. Remember what they are because I am going to ask you to name them again in a few minutes.

Please repeat these words for me: APPLE, TABLE, PENNY (may repeat names 3 times if necessary, repetition not scored)

Did the patient correctly repeat all three words?	Yes	No
1. What is the date? (D)	Correct	Incorrect
2. Name as many animals as you can in 30-seconds. (A)	_____	(number)
3. Spell "world" backwards (S)	Number Correct	0 1 2 3 4 5
4. Three-item recall (R)	Number Correct	0 1 2 3 4 5

$$\text{Brief Alzheimer's Screen} = (3.03 \times R) + (0.67 \times A) + (4.75 \times D) + (2.01 \times S)$$

BAS \leq 26 is consistent with dementia

FIGURE 5: AD8

If the patient has an accompanying reliable informant, they are asked the following questions.

Has this patient displayed any of the following issues? Remember a "yes" response indicates that you think there has been a **change in the last several years** caused by thinking and memory (cognitive) problems.

1. Problems with judgment (for example, falls for scams, bad financial decisions, buys gifts inappropriate for recipients)?
2. Reduced interest in hobbies/activities?
3. Repeats questions, stories, or statements?
4. Trouble learning how to use a tool, appliance, or gadget (VCR, computer, microwave, remote control)?
5. Forgets correct month or year?
6. Difficult handling complicated financial affairs (for example, balancing checkbook, income taxes, paying bills)?
7. Difficulty remembering appointments?
8. Consistent problems with thinking and/or memory?

Each affirmative response is one-point. A score of ≥ 2 is considered high-risk for cognitive impairment.

yes/no questions related to the clinical effects of dementia on daily life.⁴² ED personnel screening for dementia can administer the AD8 to either a patient or their caregiver (cAD8), which can be useful for non-communicative or critically ill ED patients.

Using Bayesian logic, healthcare providers can estimate the probability of dementia for these patients.^{43,44} What is Bayesian logic and how exactly is this computation accomplished in a busy ED? One needs only a pre-test probability estimate and the diagnostic accuracy (sensitivity and specificity or likelihood ratios) for the screening test. ED research indicates that the pre-test probability of dementia in patients 65 years

or older is approximately 35%.^{8, 16, 38} Next, using the ED-based diagnostic accuracy estimates from Table 1, evidence-based clinicians can derive estimates for the presence of dementia. For example, assume that the Brief Alzheimer's Score is 12, which is abnormal with a positive likelihood ratio of 2.0. One can use readily available websites (example <http://www.dokterrutten.nl/collega/LRcalcul.html>) as depicted in Figure 6. These computer-based resources are particularly useful to bypass the onerous mathematics underlying Bayesian analyses. We demonstrate the computations underlying Bayesian quantification of post-test probability in Figure 7. The Brief Alzheimer's Score of 12 yields a

FIGURE 6: Example of Online Bayesian Calculator (<http://www.dokterrutten.nl/collega/LRcalcul.html>)

Bayesian Calculator

The bayesian formula (posterior odds = LR x prior odds) is awkward in daily use, because we prefer chances over odds.

With a computer it is very easy to do the necessary calculations. In the Flash movie below the formula is integrated.

You just have to give the prior chance, like 20 (%), and the LR, like 5.5, and click the button with your mouse.

Bayesian calculations
use positive numbers

Prior chance = %

Likelihood Ratio =

Click button to calculate posterior chance  = 51.8 %

© Lex Rutten

FIGURE 7: Mathematics of Bayesian Analysis

Pre-test probability = 35%
Positive likelihood ratio = 2.2

Step 1: Convert Pre-test Probability to Pre-test Odds

$$\text{Odds} = [\text{Probability}/(1-\text{Probability})] = [0.35/(1-0.35)] = 0.35/0.65 = 0.538$$

Step 2: Multiple Pre-test Odds by Likelihood Ratio to Obtain Post-test Odds

$$\text{Post-test Odds} = \text{Pre-test Odds} * \text{Likelihood Ratio} = 0.538 * 2.2 = 1.184$$

Step 3: Convert Post-test Odds to Post-test Probability

$$\text{Probability} = [\text{Odds}/(\text{Odds}+1)] = [1.184/(1+1.184)] = 0.54$$

52% probability of dementia. On the other hand, using the same principles a Brief Alzheimer's Score ≤ 4 (negative likelihood ratio 0.10) produces a post-test probability for dementia of about 4%. Excel-based calculators are provided with the electronic version of this chapter to perform these computations for ED screeners and bedside clinicians. The AD8 is now available on iTunes as a fee application. In addition, fee-based tablet- and smartphone-based platforms like PEPID® (<http://www.pepid.com/>) now provide these dementia-screening calculators to clinicians in easy-to-use formats.

Note two key facts about the Bayesian process in screening for dementia in the ED. First, none of these dementia-screening instruments is diagnostic. A diagnostic instrument would yield post-test probabilities for dementia nearing 0% (to rule-out) and 100% (to rule-in). In particular, these screening instruments are inadequate to significantly increase the probability of dementia with positive likelihood ratios ranging 2.0-2.7. Emergency

physicians cannot diagnose or exclude dementia with these instruments. Instead, they can only provide direction for patients as to whether more time-consuming and definitive testing is indicated. Second, we do not know the exact accuracy for any of these instruments (or for most diagnostic tests in medicine). Each estimate from the research setting is accompanied by confidence intervals in **Table 1**. These confidence intervals should be considered when evaluating the merits of a diagnostic test. Specifically, would the clinical interpretation and associated actions differ significantly if the true diagnostic accuracy were one extreme of the confidence interval? If so, the diagnostic test is probably less useful. For example, the Brief Alzheimer's Screen's 95% confidence interval is 1.7 to 2.2 so the post-test probability for dementia is somewhere between 48% and 54%, although our best guess is 52% based upon the point estimate for the test's diagnostic accuracy.

Responding to Abnormal Dementia Screening Results in the Rural ED

Awareness of disease alone does not equate to better outcomes in any setting. The most pertinent challenge of using dementia-screening results in rural emergency medicine is what to do with the information. Since the screening test interpretation is not definitive, simply informing patients may produce undue anxiety without promoting recovery from the illness that brought them to the ED. Similarly, notifying caregivers may also yield unwarranted stress. A paucity of evidence exists to guide rural practice in general and dementia care is no exception.^{45,46} However, some research indicates that patients and their family value screening.⁴⁷

Emergency clinicians in rural and austere practice-settings face another challenge in managing patients with abnormal dementia screening test results. Dementia and memory service providers must address the stigma related to this diagnosis, particularly in rural areas.⁴⁸ Healthcare providers in rural settings have identified inadequate access to consultants, as well as insufficient community support and educational resources as barriers to routine dementia screening.⁴⁹ Local leaders need to target pre-hospital and hospital (nurse, emergency/primary care/specialty clinicians, and administration) leadership and outpatient resources in order to develop reliable dementia care protocols. Nursing leadership is essential to building sustainable dementia care programs within a community.^{50,51} Although administrators recognize the demographic imperative of dementia within the context of a rapidly aging society, locally pertinent financial data will be needed to propel rural-based dementia care initiatives forward.⁵¹ Primary care clinicians must be included within dementia care protocols,⁵² but the most readily identified rural barrier to standard dementia testing and ongoing care is lack of access to dementia specialists.⁵³

Dementia diagnosis and ongoing management is a collaborative effort that transcends medical specialization and compartmentalized care.⁵⁴ Fortunately, evolving technology is providing rural patients and practitioners with increased access to sub-specialty care. Specifically, telemedicine is demonstrating an increasing role in knowledge translation for a variety of high-intensity acute medicine evaluations.⁵⁵ Rural telemedicine dementia “clinics” are mutually acceptable to patients and healthcare providers, yielding enhanced clinical decision-making, more timely access to specialty care with fewer visit cancellations, and better follow-up for complex patients.⁵⁶⁻⁵⁸ Evaluating local dementia service availability as well as collaborative continuing medical education resources for nurse, physician assistant, and physician staff also promotes more rapid uptake of academic research knowledge.⁵⁹ Successful in-person^{60, 61} or remote videoconference⁶² dementia training workshops have been described.

Unanswered Questions and Future Directions

Multiple uncertainties exist about the optimal emergency care of aging adults with many opportunities for rural investigators to explore diagnostic and therapeutic models of care.^{63,64} Questions related to dementia care in the rural environment include

- Identification of the most feasible, diagnostically accurate and reliable dementia-screening instrument;
- Reproducibility of ED cognitive screening test results (are acutely ill patients less apt to display their baseline cognitive status?);
- Pragmatic screening personnel and strategies in the midst of increasing ED volumes and declining reimbursements;
- Patient and caregiver screening test result communication strategies to facilitate shared decision making;
- Efficacy and cost-effectiveness of dementia screening in improving ED chief complaint diagnostic accuracy, length of stay, and readmission rates.

Vignette Resolution

Working through the algorithm in Figure 1, your patient has an abnormal Ottawa 3DY screening test so you proceed to a formal delirium screen (see Chapter 80) which is “negative” for acute delirium. Lacking any history or physical exam evidence of trauma, you do not pursue a head CT but initiate an evaluation for acute coronary syndrome in the ED. The initial EKG is unchanged and non-diagnostic, and the first troponin is non-elevated. He displays no fluctuation in mental status during the three-hour course of care in your ED. He denies any chest pain while in the ED and his blood pressure is well-controlled, so after prescribing aspirin you call his primary care provider recommending admission to the hospital for provocative cardiac testing. While on the telephone with the primary care physician, you report the abnormal Ottawa 3DY screening test result and discuss further confirmatory testing, either in the hospital or after discharge. The primary care physician explains that she typically does not test for dementia because her rural clinic does not have access to dementia specialty care and no effective therapeutic strategies exist in 2014 anyway. You note that a rapidly expanding network of telemedicine services exist to provide rural dementia subspecialty care and the primary care physician states that she will explore those resources. You inform the patient and his family of the cardiac and cognitive test results, as well as the need for more definitive testing for dementia and the availability of telemedicine resources to expedite these work-ups. After answering their questions, you receive thanks for your thorough geriatric emergency care and resource awareness within your healthcare setting.

References

1. Pines JM, Mullins PM, Cooper JK, Feng LB, Roth KE. National trends in emergency department use, care patterns, and quality of care of older adults in the United States. *J Am Geriatr Soc.* Jan 2013;61(1):12-17.
2. Carpenter CR, Platts-Mills TF. Evolving prehospital, emergency department, and "inpatient" management models for geriatric emergencies. *Clin Geriatr Med.* Feb 2013;29(1):31-47.
3. Carpenter CR, Griffey RT, Stark S, Coopersmith CM, Gage BF. Physician and Nurse Acceptance of Geriatric Technicians to Screen for Geriatric Syndromes in the Emergency Department. *West J Emerg Med.* Dec 2011;12(4):489-495.
4. Edwards DF, Morris JC. Alone and confused: Community-residing older African Americans with dementia. *Dementia.* 2007;6(4):489-506.
5. Lewis LM, Miller DK, Morley JE, Nork MJ, Lasater LC. Unrecognized delirium in ED geriatric patients. *Am J Emerg Med.* March 1995;13(2):142-145.
6. Naughton BJ, Moran MB, Kadah H, Heman-Ackah Y, Longano J. Delirium and other cognitive impairment in older adults in an emergency department. *Ann Emerg Med.* June 1995;25(6):751-755.
7. Hustey FM, Meldon SW. The prevalence and documentation of impaired mental status in elderly emergency department patients. *Ann Emerg Med.* March 2002;39(3):248-253.
8. Carpenter CR, DesPain B, Keeling TK, Shah M, Rothenberger M. The Six-Item Screener and AD8 for the detection of cognitive impairment in geriatric emergency department patients. *Ann Emerg Med.* Jun 2011;57(6):653-661.
9. Callahan CM, Hendrie HC, Tierney MC. Documentation and evaluation of cognitive impairment in elderly primary care patients. *Ann Intern Med.* March 15 1995;122(6):422-429.
10. Valcour VG, Masaki KH, Curb JD, Blanchette PL. The detection of dementia in the primary care setting. *Arch Intern Med.* October 23 2000;160(19):2964-2968.
11. Kawas CH. Early Alzheimer's Disease. *N Engl J Med.* September 11 2003;349(11):1056-1063.
12. Blass DM, Rabins PV. In the clinic. *Dementia.* *Ann Intern Med.* April 1 2008;148(7):ITC4-1 -- ITC4-16.
13. Boustani M. Dementia screening in primary care: not too fast! *J Am Geriatr Soc.* Jul 2013;61(7):1205-1207.
14. Fox C, Lafortune L, Boustani M, Brayne C. The pros and cons of early diagnosis in dementia. *Br J Gen Pract.* Jul 2013;63(612):e510-e512.
15. Fox C, Lafortune L, Boustani M, Denning T, Rait G, Brayne C. Screening for dementia - is it a no brainer? *Int J Clin Pract.* Nov 2013;67(11):1076-1080.
16. Carpenter CR, Bassett ER, Fischer GM, Shirshekan J, Galvin JE, Morris JC. Four sensitive screening tools to detect cognitive impairment in geriatric emergency department patients: Brief Alzheimer's Screen, Short Blessed Test, Ottawa3DY, and the Caregiver Administered AD8. *Acad Emerg Med.* Apr 2011 18(4):374-384.
17. Chen R, Hu Z, Chen RL, Ma Y, Zhang D, Wilson K. Determinants for undetected dementia and late-life depression. *Br J Psychiatry.* Sep 2013;203:203-208.
18. Jia J, Wang F, Wei C, et al. The prevalence of dementia in urban and rural areas of China. *Alzheimers Dement.* 2013 (in press).
19. Forbes DA, Morgan D, Janzen BL. Rural and urban Canadians with dementia: use of health care services. *Can J Aging.* Fall 2006;25(3):321-330.
20. Lacny C, Kirk A, Morgan DG, Karunanayake C. Predictors of cognitive impairment severity in rural patients at a memory clinic. *Can J Neurol Sci.* Nov 2012;39(6):774-781.
21. Lindeman MA, Taylor KA, Kuipers P, Stothers K, Piper K. 'We don't have anyone with dementia here': a case for better intersectoral collaboration for remote Indigenous clients with dementia. *Aust J Rural Health.* Aug 2012;20(4):190-194.
22. Greenway-Crombie A, Snow P, Disler P, Davis S, Pond D. Influence of rurality on diagnosing dementia in Australian general practice. *Aust J Prim Health.* 2012;18(3):178-184.
23. Koller D, Eisele M, Kaduszkiewicz H, et al. Ambulatory health services utilization in patients with dementia - Is there an urban-rural difference? *Int J Health Geogr.* Nov 17 2010;9:59.
24. Kang Y, Meng H, Miller NA. Rurality and nursing home quality: evidence from the 2004 National Nursing Home Survey. *Gerontologist.* Dec 2011;51(6):761-773.
25. Wimo A, Winblad B, Jonsson L. The worldwide societal costs of dementia: Estimates for 2009. *Alzheimers Dement.* Mar 2010;6(2):98-103.
26. Alzheimer's Association. <http://www.alz.org/index.asp>. Accessed Dec 4, 2009.
27. Leslie DL, Marcantonio ER, Zhang Y, Leo-Summers L, Inouye SK. One-year health care costs associated with delirium in the elderly population. *Arch Intern Med.* January 14 2008;168(1):27-32.
28. Coduras A, Rabasa I, Frank A, et al. Prospective one-year cost-of-illness study in a cohort of patients with dementia of Alzheimer's disease type in Spain: The ECO Study. *J Alzheimers Dis.* Nov 5 (epub) 2010;19(2):601-615.
29. Brookmeyer R, Johnson E, Ziegler-Graham K, Arrighi HM. Forecasting the global burden of Alzheimer's disease. *Alzheimer Dement.* July 2007;3(3):186-191.
30. McCusker J, Healey E, Bellavance F, Connolly B. Predictors of repeat emergency department visits by elders. *Acad Emerg Med.* June 1997;4(6):581-588.
31. Saravay SM, Kaplowitz M, Kurek J, et al. How do delirium and dementia increase length of stay of elderly general medical inpatients? *Psychosomatics.* May-June 2004;45(3):235-242.
32. Moons P, De Ridder K, Geyskens K, et al. Screening for risk of readmission of patients aged 65 years and above after discharge from the emergency department: predictive value of four instruments. *Eur J Emerg Med.* 2007;14(6):315-323.
33. Sandhu H, Carpenter C. Clinical Decisionmaking: Opening the black box of cognitive reasoning. *Ann Emerg Med.* December 2006;48(6):713-722.
34. Folstein MF, Folstein SE, McHugh PR. "Mini-mental State": A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* November 1975;12(3):189-198.
35. Nasreddine ZS, Phillips NA, Bédirian V, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc.* Apr 2005;53(4):695-699.
36. Prince M, Acosta D, Ferri CP, et al. A brief dementia screener suitable for use by non-specialists in resource poor settings--the cross-cultural derivation and validation of the brief Community Screening Instrument for Dementia. *Int J Geriatr Psychiatry.* Sep 2011;26(9):899-907.
37. Raina SK, Raina S, Chander V, Grover A, Singh S, Bhardwaj A. Development of a cognitive screening instrument for tribal elderly population of Himalayan region in northern India. *J Neurosci Rural Pract.* Apr 2013;4(2):147-153.
38. Carpenter CR. Does this patient have dementia? *Ann Emerg Med.* Nov 2008;52(5):554-556.
39. Molnar FJ, Wells GA, McDowell I. The derivation and validation of the Ottawa 3D and Ottawa 3DY three- and four-question screens for cognitive impairment. *Clin Med Geriatrics.* 2008;2:1-11.
40. Katzman R, Brown T, Fuld P. Validation of a short Orientation-Memory-Concentration test of cognitive impairment. *Am J Psychiatry.* 1983;140:734-739.
41. Mendiondo MS, Ashford JW, Kryscio RJ, Schmitt FA. Designing a brief alzheimer screen (BAS). *J Alzheimers Dis.* October 2003;5(5):391-398.
42. Galvin JE, Roe CM, Xiong C, Morris JC. Validity and reliability of the AD8 informant interview in dementia. *Neurology.* Dec 12 2006;67(11):1942-1948.
43. Hayden SR, Brown MD. Likelihood ratio: A powerful tool for incorporating the results of a diagnostic test into clinical decisionmaking. *Ann Emerg Med.* May 1999;33(5):575-580.
44. Pines JM, Carpenter CR, Raja A, Schuur J. *Evidence-Based Emergency Care: Diagnostic Testing and Clinical Decision Rules.* 2nd ed. Oxford: Wiley-Blackwell Publishing; 2013.
45. Parsons JE, Merlin TL, Taylor JE, Wilkinson D, Hiller JE. Evidence-based practice in rural and remote clinical practice: where is the evidence? *Aust J Rural Health.* Oct 2003;11(5):242-248.

46. Innes A, Morgan D, Kosteniuk J. Dementia care in rural and remote settings: a systematic review of informal/family caregiving. *Maturitas*. Jan 2011;68(1):34-46.
47. Carpenter BD, Xiong C, Porensky EK, et al. Reaction to a dementia diagnosis in individuals with Alzheimer's disease and mild cognitive impairment. *J Am Geriatr Soc*. March 2008;56(3):405-512.
48. Szymczynska P, Innes A, Mason A, Stark C. A review of diagnostic process and postdiagnostic support for people with dementia in rural areas. *J Prim Care Community Health*. Oct 2011;2(4):262-276.
49. Teel CS. Rural practitioners' experiences in dementia diagnosis and treatment. *Aging Ment Health*. Sep 2004;8(5):422-429.
50. Morgan DG, Stewart NJ, Crossley NJ, et al. Dementia care in rural and remote areas: the first year of a CIHR new emerging team. *Can J Nurs Res*. Mar 2005;37(1):177-182.
51. Morgan D, Innes A, Kosteniuk J. Dementia care in rural and remote settings: a systematic review of formal or paid care. *Maturitas*. Jan 2011;68(1):17-33.
52. Kosteniuk J, Morgan D, Innes A, et al. Who steers the ship? Rural family physicians' views on collaborative care models for patients with dementia. *Prim Health Care Res Dev*. Apr 2013;3:1-7.
53. Bail K, Hudson C, Grealish L, et al. Characteristics of rural hospital services for people with dementia: findings from the Hospital Dementia Services Project. *Aust J Rural Health*. Aug 2013;21(4):208-215.
54. Callahan CM, Boustani MA, Weiner M, et al. Implementing dementia care models in primary care settings: The Aging Brain Care Medical Home. *Aging Ment Health*. Jan 2011;15(1):5-12.
55. Shah MN, Gillespie SM, Wood N, et al. High-intensity telemedicine-enhanced acute care for older adults: an innovative healthcare delivery model. *J Am Geriatr Soc*. Nov 2013;61(11):2000-2007.
56. Loh PK, Donaldson M, Flicker L, Maher S, Goldswain P. Development of a telemedicine protocol for the diagnosis of Alzheimer's disease. *J Telemed Telecare*. 2007;13(2):90-94.
57. Barton C, Morris R, Rothlind J, Yaffe K. Video-telemedicine in a memory disorders clinic: evaluation and management of rural elders with cognitive impairment. *Telemed J E Health*. Dec 2011;17(10):789-793.
58. Azad N, Amos S, Milne K, Power B. Telemedicine in a rural memory disorder clinic-remote management of patients with dementia. *Can Geriatr J*. Dec 2012;15(4):96-100.
59. Stark C, Innes A, Szymczynska P, Forrest L, Proctor K. Dementia knowledge transfer project in a rural area. *Rural Remote Health*. 2013;13:2060.
60. Szymczynska P, Innes A. Evaluation of a dementia training workshop for health and social care staff in rural Scotland. *Rural Remote Health*. 2011;11(2):1611.
61. Ellis IK, Philip T. Improving the skills of rural and remote generalists to manage mental health emergencies. *Rural Remote Health*. Jul-Sep 2010;10(3):1503.
62. Sclater K, Alagiakrishnan K, Sclater A. An investigation of videoconferenced geriatric medicine grand rounds in Alberta. *J Telemed Telecare*. 2004;10(2):104-107.
63. Carpenter CR, Heard K, Wilber ST, et al. Research priorities for high-quality geriatric emergency care: medication management, screening, and prevention and functional assessment. *Acad Emerg Med*. Jun 2011;18(6):644-654.
64. Carpenter CR, Shah MN, Hustey FM, Heard K, Miller DK. High yield research opportunities in geriatric emergency medicine research: prehospital care, delirium, adverse drug events, and falls. *J Gerontol Med Sci*. Jul 2011;66(7):775-783.